



Not actual patient.

Restylane
EXPRESS YOUR SMILE

Not actual patient.

- Purchase 22 Syringes of Restylane[®], Restylane-L, Restylane[®] Silk, Restylane[®] Refyne, and/or Restylane[®] Defyne and receive 22 Smile Starter Packs (\$59 Retail Value)*
- Smile Starter Packs include 1 Colorescience[®] Lip Shine SPF 35, 1 LIPSMART[®] Moisturizer

*Offer terms and conditions apply. See details below.

Promotion Offer Sheet

- “Qualifying HCPs” are all Galderma accounts with an active McKesson ID
- “Qualifying Purchase” is a purchase of 22 syringes of any combination of *Restylane-L*® 1 mL, *Restylane-L*® 0.5 mL, *Restylane*® 1 mL, *Restylane*® Silk, *Restylane*® Refyne, and/or *Restylane*® Defyne by a Qualifying HCP on a single order placed between May 6, 2019 and May 24, 2019 and delivered via a single shipment
- A “Package” includes the following Products: 1 Colorescience Lip Shine SPF 35, 1 *Lipsmart*® Moisturizer
- For each Qualifying Purchase by a Qualifying HCP, 22 Packages shall be shipped to the Qualifying HCP
- Limited quantity offer – only 66 Packages per Qualifying HCP; maximum of 10,010 total Packages for the promotion
- All orders must be placed through Galderma’s authorized distributor, McKesson Specialty Care Distribution Corporation
- This offer cannot be combined with any other offer except the ASPIRE Galderma Rewards HCP loyalty program
- Offer valid only in the U.S., excluding territories, and void where prohibited, taxed or otherwise restricted by law.
- Galderma, Colorescience, and Lipsmart each reserve the right to cancel or modify this offer without notice

Healthcare provider acknowledges that any discount or rebate provided hereunder is commensurate with the fair market value of the products. No discount or rebate provided hereunder is intended to be, nor shall it be construed as, an offer or payment made, whether directly or indirectly, to induce the referral of patients; the purchase, lease or order of any item or service from Galderma; or the recommending or arranging for the purchase, lease or order of any item or service from Galderma.

Important Safety Information

The *Restylane* family of products includes *Restylane*®, *Restylane-L*®, *Restylane*® Lyft with Lidocaine, *Restylane*® Silk, *Restylane*® Refyne, and *Restylane*® Defyne.

APPROVED USES

Restylane® and *Restylane-L*® are for mid-to-deep injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds. *Restylane*® and *Restylane-L*® are also indicated for injection into the lips in patients over the age of 21.

Restylane® Lyft with Lidocaine is for deep implantation into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds and for cheek augmentation and for the correction of age-related midface contour deficiencies in patients over the age of 21. *Restylane*® Lyft with Lidocaine is also indicated for injection into the subcutaneous plane in the dorsal hand to correct volume deficit in patients over the age of 21.

Restylane® Silk is for lip augmentation and for correction of perioral wrinkles in patients over the age of 21.

Restylane® Refyne is for mid-to-deep injection into the facial tissue for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds, in patients over the age of 21.

Restylane® Defyne is for mid-to-deep injection into the facial tissue for the correction of moderate to severe deep facial wrinkles and folds, such as nasolabial folds, in patients over the age of 21.

Are there any reasons why I should not use products within the Restylane® family? (Contraindications)

To ensure a safe procedure, your doctor will talk to you about your medical history to determine if you are an appropriate candidate for treatment. You should not use products within the Restylane family if:

- You have severe allergies with a history of severe reactions (anaphylaxis)

Important Safety Information, continued

- You are allergic to lidocaine or to any of the gram-positive bacterial proteins used to make hyaluronic acid
- You are prone to bleeding or have been diagnosed with a bleeding disorder

Are there other precautions that I should discuss with my doctor?

- Tell your doctor if you are breastfeeding, pregnant, or trying to become pregnant. The safety of these products for use during pregnancy, or in women who are breastfeeding, has not been studied
- *Restylane*, *Restylane-L*, *Restylane*® Lyft with Lidocaine, *Restylane* Refyne and *Restylane* Defyne are intended to treat facial wrinkles and folds, such as nasolabial folds. *Restylane* and *Restylane-L* are also intended for lip enhancement. *Restylane*® Lyft with Lidocaine is also intended for injection in the dorsal hand to correct volume loss. Treatments in other areas of the face or body have not been evaluated in clinical studies.
- The safety and effectiveness of *Restylane*® Silk for areas other than the lips and perioral area have not been evaluated in clinical studies.
- Tell your doctor if you have any history of scarring, particularly thick and stiff scars, or any pigmentation (skin color) disorders. These side effects can occur with hyaluronic acid fillers in general.
- Tell your doctor if you are planning other laser treatments or a chemical peel, as there is a possible risk of inflammation at the treatment site if these procedures are performed after treatment
- Patients who experience skin injury near the site of injection with these products may be at a higher risk for side effects
- Tell your doctor if you are on any medications to decrease your body’s immune response (immunosuppressive therapy). Using these medications may increase your risk of bruising or bleeding at the gel injection site.
- Tell your doctor if you are using any “blood thinners” such as aspirin, warfarin, or any other medications that affect bleeding. Using these medications may increase your risk of bruising or bleeding at the gel injection site.
- The use of these products on gel injection sites with skin sores, pimples, rashes, hives, cysts, or infections should be postponed until healing is complete. Use of product in these areas could delay healing or make your skin problems worse.
- Tell your doctor if you have diseases, injuries, or disabilities of the hand.

What are the possible side effects?

The most commonly observed side effects are swelling, redness, pain, bruising, headache, tenderness, lump formation, itching at the injection site, and impaired hand function. These are typically mild in severity and typically resolve in less than 7 days in nasolabial folds and less than 14 days in lips. Serious but rare side effects include delayed onset infections, recurrence of herpetic eruptions, and superficial necrosis at the injection site.

One of the risks with using this product is unintentional injection into a blood vessel. The chances of this happening are very small, but if it does happen, the complications can be serious, and may be permanent. These complications, which have been reported for facial injections, can include vision abnormalities, blindness, stroke, temporary scabs, or permanent scarring of the skin.

As with all skin injection procedures, there is a risk of infection.

To report a side effect with any of the Restylane products, please call Galderma Laboratories, L.P. at 1-855-425-8722.

The *Restylane* family of products is available only through a licensed practitioner. Complete Instructions for Use are available at www.RestylaneUSA.com.